

WHAT IS CLAIMED IS:

1. An isolated DNA sequence encoding a 30-kilodalton protein of *Ehrlichia canis*, wherein said protein is immunoreactive with anti-*Ehrlichia canis* serum.

2. The DNA sequence of claim 1, wherein said protein comprises of an amino acid sequence selected from the group consisting of SEQ ID No. 2, 4, 6, 40, 42, 44 and 46.

3. The DNA sequence of claim 2, wherein said protein has an N-terminal signal sequence.

4. The DNA sequence of claim 3, wherein said protein is post-translationally modified to a 28-kilodalton protein.

5. The DNA sequence of claim 1, wherein said DNA comprises a sequence selected from the group consisting of SEQ ID No. 1, 3, 5, 39, 41, 43 and 45.

5

6. The DNA sequence of claim 1, wherein said DNA is contained in a single locus of *Ehrlichia canis*.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

7. The DNA sequence of claim 6, wherein said locus is a multigene locus of 10,677 base pairs in length.

8. The DNA sequence of claim 7, wherein said locus contains genes encoding homologous 28-kilodalton proteins of *Ehrlichia canis*.

9. The DNA sequence of claim 8, wherein said homologous 28-kilodalton proteins of *Ehrlichia canis* are selected

from the group consisting of p28-1, p28-2, p28-3, p28-4, p28-5, p28-6, p28-7, p28-8 and p28-9.

5 10. A vector comprising the DNA sequence of claim 1.

10 11. The vector of claim 10, wherein said vector is an expression vector capable of expressing a peptide or polypeptide encoded by a sequence selected from the group consisting of SEQ ID No. 1, 3, 5, 39, 41, 43 and 45 when said expression vector is introduced into a cell.

15 12. A recombinant protein comprises of an amino acid sequence selected from the group consisting of SEQ ID No. 2, 4, 6, 40, 42, 44 and 46.

20 13. The recombinant protein of claim 12, wherein said amino acid sequence is encoded by a nucleic acid segment

comprising a sequence selected from the group consisting of SEQ ID No. 1, 3, 5, 39, 41, 43 and 45.

5 14. A host cell comprising a nucleic acid segment selected from the group consisting of SEQ ID No. 1, 3, 5, 39, 41, 43 and 45.

10 15. A method of producing the recombinant protein of claim 12, comprising the steps of:

obtaining a vector that comprises an expression region comprising a sequence encoding the amino acid sequence selected from the group consisting of SEQ ID No. 2, 4, 6, 40, 42, 44 and 46
15 operatively linked to a promoter;

transfecting said vector into a cell; and

culturing said cell under conditions effective for expression of said expression region.

20

16. An antibody immunoreactive with a polypeptide comprises of an amino acid sequence selected from the group consisting of SEQ ID No. 2, 4, 6, 40, 42, 44 and 46.

5

17. A method of inhibiting *Ehrlichia canis* infection in a subject comprising the steps of:

identifying a subject prior to exposure or suspected of being exposed to or infected with *Ehrlichia canis*; and

10 administering a composition comprising a 28-kDa antigen of *Ehrlichia canis* in an amount effective to inhibit *Ehrlichia canis* infection.

15

18. The method of claim 17, wherein said 28-kDa antigen is a recombinant protein comprising an amino acid sequence selected from the group consisting of SEQ ID No. 2, 4, 6, 40, 42, 44 and 46.

20

[illegible]

28-kDa and carrier.